

Inertia-gravity waves and geostrophic turbulence

William R. Young[†]

Scripps Institution of Oceanography, University of California San Diego, La Jolla, CA 92093-0213, USA

(accepted 12 April 2021)

Inertia-gravity waves in the atmosphere and ocean are transported and refracted by geostrophic turbulent currents. Provided that the wave group velocity is much greater than the speed of geostrophic turbulent currents, kinetic theory can be used to obtain a comprehensive statistical description of the resulting interaction (Savva *et al.*, *J. Fluid Mech.*, vol. 916, 2021, A6). The leading-order process is scattering of wave energy along a surface of constant frequency, ω , in wavenumber space. The constant- ω surface corresponding to the linear dispersion relation of inertia-gravity waves is a cone extending to arbitrarily high wavenumbers. Thus, wave scattering by geostrophic turbulence results in a cascade of wave energy to high wavenumbers on the surface of the constant- ω cone. Solution of the kinetic equations shows establishment of a wave kinetic energy spectrum $\sim k_h^{-2}$, where k_h is the horizontal wavenumber.

Key words: atmospheric flows, oceanography, geostrophic turbulence

1. Introduction and background

Within the rapidly rotating fluid envelope of the Earth, slow geostrophic turbulence co-exists – not entirely peacefully – with fast inertia-gravity waves (hereafter IGWs): see [figure 1](#). Geostrophic turbulence refers to a form of two-dimensional turbulence with geophysical complications arising from density stratification and the planetary β -effect (Charney 1971). A main departure of geostrophic turbulence from plain and simple two-dimensional turbulence is that horizontal velocities are vertically sheared. Weather systems in the atmosphere, with an evolutionary time scale of several days, are a familiar example of geostrophic turbulence. The IGWs, also known as internal waves, are higher-frequency motions that propagate vertically in a stably stratified fluid and involve a balance between inertia, buoyancy, pressure gradient and Coriolis forces. The IGW time scales are of the order of hours i.e. much shorter than those of geostrophic turbulence.

The interaction between waves and turbulence, with widely separated time scales, presents meteorologists and oceanographers with a ‘wave–turbulence jigsaw’ (McIntyre 2008). Within the last decade, several pieces of this sprawling puzzle have quietly dropped into place. This advance has greatly clarified the extent to which there is a separation in

[†] Email address for correspondence: wryoung@ucsd.edu




Figure 1. Co-existence of geostrophic turbulence and IGWs. The figure shows horizontal slices through a three-dimensional solution of the Boussinesq equations. The geostrophic turbulence in the lower panel is visualized by showing vertical vorticity; the IGWs in the upper panel are revealed with vertical velocity. Figure contributed by H. Kafiabad.

length scale between geostrophic turbulence and IGWs. The short answer is that there is not as much separation in length scale as meteorologists and oceanographers expected: IGWs are energetic on surprisingly large scales. Geophysical kinetic energy spectra have a band of wavenumbers within which waves and turbulence are equally energetic. A painful consequence of this length-scale overlap is that the approximation of Wentzel, Kramers and Brillouin (WKB hereafter) does not apply to all scales of interest.

Separating waves from turbulence in geophysical energy spectra has required new statistical tools for the analysis of observations of fluid velocity along the one-dimensional transects made by ships and planes (Bühler, Callies & Ferrari 2014). Aircraft data show that atmospheric geostrophic turbulence dominates the synoptic range, while IGWs dominate the mesoscale range (Callies, Ferrari & Bühler 2014; Waite 2020). The transition scale between the two ranges is at around 500 km. The oceanic situation is more complicated because the ocean is more spatially inhomogeneous than the atmosphere, and because geostrophic eddies in the ocean are much smaller than their atmospheric cousins. But the broad-brush conclusion is the same: ocean IGWs are energetic on surprisingly large length scales. For instance, Rocha *et al.* (2016) show that in Drake Passage, with a deformation radius of 16 km, IGWs account for roughly half of the near-surface kinetic energy at scales between 10 and 40 km.

Synthesis of oceanographic IGW data into a seemingly ‘universal spectrum’ (Garrett & Munk 1972) drove intensive research on nonlinear wave interactions in the seventies and eighties (Müller *et al.* 1986). While this effort did not ignore the interaction of IGWs with geostrophic turbulence – see for instance Müller (1976) – the focus was mainly on self-interactions within the IGW field as an explanation of the Garrett–Munk spectrum. A development driving a re-examination of the IGW spectrum is the realization that geostrophic turbulence is the main reservoir of ocean kinetic energy (Ferrari & Wunsch 2009). The interaction of IGWs with this turbulent reservoir is likely to be an important mechanism for shaping the IGW spectrum – perhaps more important than wave–wave interactions.

2. Overview of Savva, Kafiabad & Vanneste (2021)

The recent paper by Savva *et al.* (2021, SKV hereafter) is the first comprehensive and definitive study of IGW scattering by geostrophic turbulence. The crucial assumption of

SKV is the weak-current approximation that

$$|U|/c_g \ll 1. \tag{2.1}$$

Here, $|U|$ denotes the typical speed of geostrophic currents and c_g is the typical group speed of IGWs. The condition (2.1) ensures that wave packets rapidly propagate through many decorrelation lengths of the turbulent velocity field and that during this passage the turbulence does not evolve significantly. Thus the scattering velocity field is effectively frozen, the Doppler shift is negligible, the intrinsic frequency of the waves is unchanged by the interaction and conservation of action (Bretherton & Garrett 1968) is, to leading order, the same as conservation of IGW energy. Because there is almost no transfer of energy between waves and turbulence the interaction is catalytic i.e. the wave field is modified by turbulence (details below) but the turbulence is unaffected by the waves. Thus SKV treats the turbulence as a random velocity field with a specified kinetic energy spectrum $E_K(\mathbf{k})$, where \mathbf{k} is the wavenumber.

SKV avoids the WKB approximation by using the Wigner transform formalism of Ryzhik, Papanicolaou & Keller (1996) and shows that the phase space energy density of the waves, denoted $a(\mathbf{k}, \mathbf{x}, t)$, satisfies the kinetic equation

$$\partial_t a(\mathbf{x}, \mathbf{k}, t) + \nabla_{\mathbf{k}} \omega(\mathbf{k}) \cdot \nabla_{\mathbf{x}} a(\mathbf{x}, \mathbf{k}, t) = \int \sigma(\mathbf{k}, \mathbf{k}') a(\mathbf{x}, \mathbf{k}', t) d\mathbf{k}' - \Sigma(\mathbf{k}) a(\mathbf{x}, \mathbf{k}, t), \tag{2.2}$$

where $\Sigma(\mathbf{k}) = \int \sigma(\mathbf{k}, \mathbf{k}') d\mathbf{k}'$. Above $\omega(\mathbf{k})$ is the IGW dispersion relation

$$\omega(\mathbf{k}) = \pm \sqrt{\frac{N^2 k_h^2 + f^2 k_3^2}{k_h^2 + k_3^2}}, \tag{2.3}$$

where N is the buoyancy frequency and f is the Coriolis frequency; the wavenumber is decomposed into horizontal and vertical components $\mathbf{k} = (k_h, k_3)$. The surface of constant ω in \mathbf{k} -space defined by (2.3) is a double cone.

The scattering cross-section in (2.2) has the form

$$\sigma(\mathbf{k}, \mathbf{k}') = \mathcal{E}(\mathbf{k}, \mathbf{k}') E_K(\mathbf{k}' - \mathbf{k}) \delta(\omega(\mathbf{k}') - \omega(\mathbf{k})), \tag{2.4}$$

where $\mathcal{E}(\mathbf{k}, \mathbf{k}') = \mathcal{E}(\mathbf{k}', \mathbf{k})$ is given by a formidable expression in SKV. The kinetic equation (2.2) subsumes earlier studies of special cases (Danioux & Vanneste 2016; Savva & Vanneste 2018; Kafiabad, Savva & Vanneste 2019).

The $\delta(\omega(\mathbf{k}') - \omega(\mathbf{k}))$ term in (2.4) ensures that ω is unchanged by scattering. The interaction can be viewed as a resonant triad between two IGWs and a zero-frequency geostrophic mode. The upshot is that all the IGW energy that starts on a particular constant- ω double cone stays on that same double cone. If the turbulence is horizontally isotropic then scattering of IGW energy over the surface of the \mathbf{k} -space double cone involves three processes:

- (a) The horizontal rate of strain of geostrophic turbulence results in horizontal isotropization by azimuthal scattering of IGWs around the cone, with k_h and k_3 unchanged (Savva & Vanneste 2018).
- (b) The vertical shear of geostrophic turbulence scatters IGW energy along the k_3 -axis of the cone and so increases $|\mathbf{k}|$.
- (c) Energy is weakly transferred between the two halves of the double cone via inelastic scattering (McComas & Bretherton 1977).

A main result from SKV is the cascade of IGW energy to high wavenumbers in (b). The special role of vertical shear in enabling turbulent scattering to access the entire constant ω double cone is notable. This high-wavenumber IGW cascade relies on a peculiar property of the dispersion relation (2.3): the conical constant- ω surface is not compact so that scattering with constant ω can access arbitrarily high wavenumbers (The ω -surface for acoustic scattering is a sphere in \mathbf{k} -space; a sphere is compact and thus turbulence cannot catalyse a cascade of acoustic energy to high wavenumbers).

As IGWs are scattered out to high wavenumbers on the cone, the WKB-based induced diffusion approximation of McComas & Bretherton (1977) becomes applicable and enables a great simplification of (2.2): the non-local transfers on the right-hand side are approximated by \mathbf{k} -space diffusion along the surface of the cone. Kafiabad *et al.* (2019) solve this simplified version of (2.2) with analytic methods. This solution shows that induced diffusion results in a k_h^{-2} spectrum of wave energy. Now k_h^{-2} is a frequently observed energy spectrum in the ocean; for example, Rocha *et al.* (2016) show that the IGW component of the kinetic energy spectrum is k_h^{-2} . In the atmosphere the shallow mesoscale part of the kinetic energy spectrum is traditionally described as a $k_h^{-5/3}$ spectrum (Nastrom & Gage 1985). Atmospheric data are, however, also consistent with k_h^{-2} . Kafiabad *et al.* (2019) speculate that these observations in the ocean and atmosphere could be explained by the k_h^{-2} spectrum resulting from induced diffusion of IGW energy by geostrophic turbulence.

3. Future

The unsteady evolution of geostrophic turbulence results in weak scattering of IGW energy off the constant- ω double cone. This cross ω -surface diffusion has been demonstrated by Dong, Bühler & Shafer Smith (2020) using the shallow water equations and the induced diffusion approximation. Diffusion across the ω -surfaces implies an increase in wave energy, so that IGWs act as an effective viscosity on the turbulence (Müller 1976). The next step is to investigate this effect in the Boussinesq equations and include it in the kinetic equation (2.2). In principle this can be accomplished by computing the cross- ω -surface scattering as the next term in the expansion that leads to (2.2).

Another frontier is strong multiscale interactions – but not so strong as in stratified turbulence – leading to failure of (2.1). Special examples, such as the strong wave–wave interaction discussed by Broutman & Young (1986), show large changes in frequency and significant energy transfers. This is also likely the case for strong interactions between IGWs and geostrophic turbulence. Provided that there is a separation in length scales, this frontier problem seems approachable only via WKB and Monte Carlo simulation.

Acknowledgements. Thanks to O. Bühler, P. Cessi, H. Kafiabad and J. Vanneste for comments and discussion.

Funding. This work was supported by the National Science Foundation (award number OCE-2048583) and by the Office of Naval Research (award number N00014-18-1-2803).

Declaration of interest. The author reports no conflict of interest.

Author ORCID.

William R. Young <https://orcid.org/0000-0002-1842-3197>.

REFERENCES

BRETHERTON, F.P. & GARRETT, C.J.R. 1968 Wavetrains in inhomogeneous moving media. *Proc. R. Soc. Lond. A Math. Phys. Sci.* **302**, 529–554.

Inertia-gravity waves and geostrophic turbulence

- BROUTMAN, D. & YOUNG, W.R. 1986 On the interaction of small-scale oceanic internal waves with near-inertial waves. *J. Fluid Mech.* **166**, 341–358.
- BÜHLER, O., CALLIES, J. & FERRARI, R. 2014 Wave–vortex decomposition of one-dimensional ship-track data. *J. Fluid Mech.* **756**, 1007–1026.
- CALLIES, J., FERRARI, R. & BÜHLER, O. 2014 Transition from geostrophic turbulence to inertia–gravity waves in the atmospheric energy spectrum. *Proc. Natl Acad. Sci.* **111**, 17033–17038.
- CHARNEY, J. 1971 Geostrophic turbulence. *J. Atmos. Sci.* **28**, 1087–1095.
- DANIOUX, E. & VANNESTE, J. 2016 Near-inertial wave scattering by random flows. *Phys. Rev. Fluids* **1**, 033701.
- DONG, W., BÜHLER, O. & SHAFER SMITH, K. 2020 Frequency diffusion of waves by unsteady flows. *J. Fluid Mech.* **905**, R3.
- FERRARI, R. & WUNSCH, C. 2009 Ocean circulation kinetic energy: reservoirs, sources, and sinks. *Annu. Rev. Fluid Mech.* **41**, 253–282.
- GARRETT, C. & MUNK, W. 1972 Space-time scales of internal waves. *Geophys. Fluid Dyn.* **3**, 225–264.
- KAFIABAD, H., SAVVA, M.A.C. & VANNESTE, J. 2019 Diffusion of inertia-gravity waves by geostrophic turbulence. *J. Fluid Mech.* **869**, R7.
- MCCOMAS, C.H. & BRETHERTON, F.P. 1977 Resonant interaction of oceanic internal waves. *J. Geophys. Res.* **82**, 1397–1412.
- MCINTYRE, M.E. 2008 Potential-vorticity inversion and the wave-turbulence jigsaw: some recent clarifications. *Adv. Geosci.* **15**, 47–56.
- MÜLLER, P. 1976 On the diffusion of momentum and mass by internal gravity waves. *J. Fluid Mech.* **77**, 789–823.
- MÜLLER, P., HOLLOWAY, G., HENYEV, F. & POMPHREY, N. 1986 Nonlinear interactions among internal gravity waves. *Rev. Geophys.* **24**, 493–536.
- NASTROM, G.D. & GAGE, K.S. 1985 A climatology of atmospheric wavenumber spectra of wind and temperature observed by commercial aircraft. *J. Atmos. Sci.* **42**, 950–960.
- ROCHA, C.B., CHERESKIN, T.K., GILLE, S.T. & MENEMENLIS, D. 2016 Mesoscale to submesoscale wavenumber spectra in Drake Passage. *J. Phys. Oceanogr.* **46**, 601–620.
- RYZHIK, L., PAPANICOLAOU, G. & KELLER, J.B. 1996 Transport equations for elastic and other waves in random media. *Wave Motion* **24**, 327–370.
- SAVVA, M.A.C., KAFIABAD, H.A. & VANNESTE, J. 2021 Inertia-gravity-wave scattering by three-dimensional geostrophic turbulence. *J. Fluid Mech.* **916**, A6.
- SAVVA, M.A.C. & VANNESTE, J. 2018 Scattering of internal tides by barotropic quasigeostrophic flows. *J. Fluid Mech.* **856**, 504–530.
- WAITE, M.L. 2020 Untangling waves and vortices in the atmospheric kinetic energy spectra. *J. Fluid Mech.* **888**, F1.